

Final

Juvenile Collection and Bypass Report, 2024

Lower Monumental Project Juvenile Fish Facility

Prepared for



US Army Corps of Engineers

Lower Monumental Dam

5520 Devil Canyon Road
Kahlotus, WA 99335

Prepared by

Environmental Assessment Services, LLC
350 Hills Street, Suite 112
Richland, Washington 99354
October 2024

Prepared by:

Matt Paulsen
Mark Morasch
Phoebe Hill
Environmental Assessment Services
350 Hills St, Suite 112
Richland, Washington 99354
15 October 2024

Table of Contents

SUMMARY.....	1
ACKNOWLEDGEMENTS.....	1
FACILITY INTRODUCTION AND DESCRIPTION.....	1
FACILITY MODIFICATIONS/MAINTENANCE AND IMPROVEMENTS	2
RIVER CONDITIONS.....	2
River Flow	2
River Temperature.....	3
JUVENILE BYPASS.....	1
Migration, Sampling, and Bypass of Juvenile Salmonids	1
Collection.....	1
Sampling.....	4
Transportation-Barge Loading Operations.....	7
Transportation-Truck Loading Operations	7
Bypass.....	7
Migration, Sampling, and Bypass of Juvenile Lamprey.....	8
Incidental Species.....	8
Adult Fallbacks.....	9
Separator Efficiency.....	10
FISH CONDITION.....	13
Descaling	13
Other Injury and Disease	16
Mortality.....	16
RESEARCH	20
Gas Bubble Trauma.....	20
PNNL Lamprey Study	20
Steelhead Kelt Study	20
FACILITY OPERATIONS AND MAINTENANCE.....	20
Turbine Operations	20
Spill Operations.....	21
Removable Spillway Weir	21
Forebay Debris	22
Trash Racks	22
Gatewells	22
Submersible Traveling Screens	22
Vertical Barrier Screens	23
Juvenile Collection Channel Orifices	23
Primary Dewatering Structure.....	23
JUVENILE FISH FACILITY	24
Separator	24
PIT-Tag System	24
Sample	24
Other Facility Issues	25
FISH SALVAGE.....	25

COOLING WATER STRAINERS	26
INVASIVE SPECIES.....	26
AVIAN PREDATION.....	27
Avian Predation-General.....	27
Adult Fishway Inspection Bird Counts/Wildlife Services Bird Monitoring	28
Gulls	28
Cormorants	29
Terns	29
Grebes	30
Pelicans.....	30
Avian Hazing-United States Department of Agriculture-Wildlife Services.....	30
RECOMMENDATIONS.....	30

List of Tables

Table 1.	Comparison of average monthly flow and spill at Lower Monumental Dam, 2019–2024 and the 5-year average.	2
Table 2.	Average monthly river temperatures (°F), 2019–2024 and 5-year average.	3
Table 3.	Annual collection, bypass, and transport at Lower Monumental Dam, 2020-2024.	2
Table 4.	Annual peak collection dates at Lower Monumental Dam, 2020-2024.	3
Table 5.	Annual percentage sampled out of total collection for each juvenile salmonid species group at Lower Monumental Dam, 2020-2024.	5
Table 6.	Weekly average sample rates in percent and sample totals at Lower Monumental Dam, 2024.	5
Table 7.	Pacific lamprey in the sample and separator, and the total collection for both life stages, with mortality 2020–2024.	8
Table 8.	Estimated collection of incidental species at Lower Monumental Dam, 2024.	9
Table 9.	Annual totals of adult salmonids released from the separator at Lower Monumental Dam, 2020–2024.	10
Table 10.	Monthly totals of adult salmonids released from the separator at Lower Monumental Dam, 2024.	10
Table 11.	Condition of adult salmonids released from the separator at Lower Monumental Dam, 2024.	10
Table 12.	Annual separator efficiency in percent at Lower Monumental Dam, 2020–2024.	12
Table 13.	Annual descaling rates in percent for fish sampled at Lower Monumental Dam, 2020–2024.	14
Table 14.	Weekly descaling rates in percent for fish sample at Lower Monumental Dam, 2024.	14
Table 15.	Annual facility mortality in percent at Lower Monumental Dam, 2020-2024.	18
Table 16.	Weekly facility mortality in percent at Lower Monumental Dam, 2024.	18
Table 17.	Annual sample mortality in percent at Lower Monumental Dam, 2020-2024.	19
Table 18.	Unit outages and causes at Lower Monumental, 2023.	21
Table 19.	Cooling Water Strainer Results at Lower Monumental Dam, 2023.	26
Table 20.	USDA-WS hazing program schedule for Lower Monumental Dam, 2023.	30

List of Figures

Figure 1.	Comparison of daily average river flow and spill at Lower Monumental Dam, 2024.	3
Figure 2.	Daily juvenile salmonid collection, all species combined, versus daily average river flow at Lower Monumental Dam, 2024.	4
Figure 3.	Daily count of gulls, cormorant, grebes and pelicans in the tailrace and forebay, 2023.	28

List of Acronyms

BiOp	Biological Opinion for Operations
BPA	Bonneville Power Administration
CFS	cubic feet per second
FPC	Fish Passage Center
GBT	gas bubble trauma
JCC	Juvenile Collection Channel
JFF	Juvenile Fish Facility
JSATS	Juvenile Salmon Acoustic Telemetry System
KCFS	kilo cubic feet per second
LMN	Lower Monumental Dam
LWG	Lower Granite Dam
ODFW	Oregon Department of Fish and Wildlife
OOS	out of service
PIT	passive integrated transponder
PDS	primary dewatering structure
PNNL	Pacific Northwest National Laboratory
STS	submersible traveling screens
RSW	removable spill weir
USACE	U.S. Army Corps of Engineers
USDA-WS	United States Department of Agriculture-Wildlife Services
VBS	vertical barrier screen

Summary

Juvenile fish transportation and bypass operations occurred at Lower Monumental Dam (LMN) Juvenile Fish Facility (JFF) in 2024. Submersible traveling screens (STSs) for all operating units were installed between March 18 and March 20. Collection channel returned to service and was watered up at 0915 on March 21. Primary dewatering structure returned to service at the same time. During collection channel watering up checks, all faulty orifice lights were replaced. The JFF returned to service and was watered up at 1015 on March 21. Sampling occurred at 0700 until April 23, when a partial sample occurred at 1300, from a collection period of 0700 to 1300, to transition to every-day sampling on a 1300 to 1300 schedule for transport season (April 24 to June 20). Collection/sampling was suspended two times due to high river temperatures during the 2024 season, on July 22, and August 20.

Fish were transported by barge from April 24 through May 4, May 18 to June 1, then every fourth day from June 1 to June 17. The fish passage plan requires fish to be transported from April 24 to June 20, but barging was halted twice in 2024 due to high flows. The total number of fish bypassed on those days was 15,746. No truck transport occurred at LMN in 2024.

Total smolt collection in the 2024 season was 159,209. This includes expanded numbers of those sampled during pre-transport. Of the fish collected in the 2024 season, 84,238 were barged, and 74,724 were bypassed, and there were 247 mortalities. This season's total collection by species group included 36,009 clipped yearling Chinook salmon, 6,146 unclipped yearling Chinook salmon, 15,813 clipped subyearling Chinook salmon, 30,414 unclipped subyearling Chinook salmon, 53,798 clipped steelhead, 11,872 unclipped steelhead, 558 unclipped sockeye/kokanee salmon, and 2,295 combined clipped/unclipped coho salmon (Table 3). A total of 2,304 Chinook salmon fry were collected during the 2024 season.

Oregon Department of Fish and Wildlife (ODFW) technicians examined 1,124 fish for gas bubble trauma (GBT) in 2024. Examinations were conducted once a week from April 7 through July 31.

Acknowledgements

USACE personnel included supervisory biologist Denise Griffith; assistant biologist Raymond Addis; biological technicians and maintenance personnel. Representing Environmental Assessment Services were Phoebe Hill, Mark Morasch, and Matt Paulsen. Technical editing was performed by Joanne Stover. Thank you to all crews from USACE and various state, federal, and tribal entities for making this a successful season.

Facility Introduction and Description

LMN is located at river mile 41.6 on the Snake River. The dam is located at the head of Lake Sacajawea, the reservoir created by Ice Harbor Dam. Lower Monumental has six 135,000-kilowatt turbine units. To bypass the turbines, the juvenile system begins with trash racks, STSs and VBSs. When fish enter each turbine intake, they are diverted into the gatewell slots by the STSs. Each unit has three gatewell slots. Each gatewell slot has two orifices where lights are directed at each open orifice to encourage fish movement into the collection channel. The fish pass through these twelve-inch orifices to the JCC which terminates at the PDS where all but 30 cubic feet second (cfs) flow is removed. The remaining 30 cfs flow and fish are routed through the transport flume to the separator. Upon reaching the separator, non-target adult fish are released to the river and juvenile fish pass below the separator bars and enter the distribution system. The full distribution system includes transport flumes, passive integrated transponder (PIT) tag bypass, sampling facilities, holding facilities, and barge and truck loading capabilities.

Downstream of the separator in the A and B flumes are the PIT-tag bypass. The A and B sides each have a set of sample and PIT-tag systems. Inside the JFF building is the wet lab where the fish condition sample occurs.

Finally, LMN has eight spillbays with a removable spillway weir (RSW) in spillbay 8. The RSW was installed to provide a surface passage route and improve conditions for out-migrating juveniles.

Facility Modifications/Maintenance and Improvements

Maintenance and improvements for the 2023 year which are made to enhance the system performance over the previous seasons include a new raceway tailscreen hoist and removal of old wet lab infrastructure and piping. The wet floor lab was also resealed on March 11.

A new STS camera was purchased in March 2023 and used throughout the season. It was hard wired to the STS van on August 22 and appeared to work perfectly throughout the year.

All flume leaks were repaired in October by removing old silicone and replacing new silicone where the old gasket material had decomposed, once all the flumes had been power washed. All cracked and leaking oil misters were also replaced in October. Approximately five of various sizes were replaced.

Also, during the fall, large logs and washed-up trash was removed from the inside fenced area of the outfall pipe pump. Both fire hydrants at the JFF were refurbished with all new parts in October.

River Conditions

River Flow

During the 2024 season, the average daily river flow did not exceed 114.9 thousand cubic feet per second (kcfs) (Figure 1). The average flow for the season was 47.8 kcfs, with the highest daily average flow for the season of 114.9 kcfs on June 5, and the lowest daily average flow for the season occurring on September 27 with a flow of 14.3 kcfs. Spring spill mandated by the National Oceanic and Atmospheric Administration Biological Opinion for Operation of the Federal Columbia River Power System (BiOp) occurred for 79 days from April 3 through midnight on June 20, with a maximum daily average spill of 94.5 kcfs on May 18. The RSW was put into operation for spring steelhead migration on March 1 to March 31 and spring spill at 0100 hours on April 3, following the initiation of BiOp-mandated spill. Summer spill ended at 2359 hours on August 31. The 2024 river flow was lower than the 5-year average during the months of May, June, July, August, and September, and flows were higher than the 5-year average during the months of March and April. In 2024, spill averages were lower than the 5-year average during the months of July and August, and spill averages were higher than the 5-year average during the months of March, April, May June, and September. Summer spill was higher than the 5-year average. Average monthly flow and spill for the 2019 to 2024 collection seasons are provided in Table 1.

Table 1. Comparison of average monthly flow and spill at Lower Monumental Dam, 2019–2024 and the 5-year average.

Month	2019	2020	2021	2022	2023	2019–23 Average	2024
Flow (kcfs)							
March	--	--	--	--	26.9	26.9	47.9
April	115.8	53.5	51	42.2	53.1	63.1	67.5
May	62.6	103.7	68.9	84	132	90.2	78.7
June	87.8	92.9	52.7	132.4	69.8	87.1	68.8
July	37.7	47.3	26.9	44.6	36.9	38.7	33.4
August	26.8	26.8	22.9	28.5	26.4	26.3	25.9
September	24.4	22.8	18.9	20	22.9	21.8	19.9
Spill (kcfs)							
March	--	--	--	--	0.6	0.6	8.0
April	41.3	35.9	33.7	24.2	36.1	34.2	52.4

May	38.3	71.7	47.9	55.2	69	56.4	67.1
June	36.4	54.9	33.1	62.2	36.4	44.6	49.0
July	16.8	17.4	13.4	17	17.1	16.3	16.0
August	13.5	11.2	9.4	11.7	11.1	11.4	8.4
September	0.2	0.2	0.2	0.8	0.8	0.4	1.5




Figure 1. Comparison of daily average river flow and spill at Lower Monumental Dam, 2024.

River Temperature

River temperature is measured daily in the JFF wetlab freshwater supply at 1300 hours during barging season, and at 0700 hours during periods of bypass. Temperatures observed in the wetlab were higher than the 5-year average for the months of March, May, June, and July. Water temperatures in the wetlab during the 2024 season were lower than the 5-year average during the months of April, August, and September (Table 2). River temperature averaged 61.8°F for the 2024 season and ranged from the low of 46.4°F on March 26, to the high of 70.3°F recorded on July 24, 25, and August 19.

Table 2. Average monthly river temperatures (°F), 2019–2024 and 5-year average.

Month*	2019	2020	2021	2022	2023	2019–23 Average	2024
April	49.3	49.9	49.4	48.6	48.0	49.0	47.4
May	54.8	53.4	55	52.3	54.8	54.1	55.6
June	61.6	58.9	61.5	55.9	62.6	60.1	60.5
July	67.1	65.8	69.6	66.6	68.6	67.5	68.1
August	69.1	68.8	69.9	69.4	69.8	69.4	69.2
September	68.0	66.4	66.4	67.7	67.1	67.1	66.5

Note:

* March and October were not included in the table, we sampled only two days in March and one in October.

Juvenile Bypass

Migration, Sampling, and Bypass of Juvenile Salmonids

Collection

Pre-transport primary bypass occurred from 0700 hours April 1 through 0700 hours April 23. Fish collection for barge transportation began at 0700 hours on April 23 and continued until 0700 hours June 17. An estimated 159,209 juvenile salmonids were collected in 2024 (Table 3). Within each species group, the number collected, and percent of the total collection was 36,009 clipped yearling Chinook salmon (22.6%), 6,146 unclipped yearling Chinook salmon (3.9%), 15,813 clipped subyearling Chinook salmon (9.3%), 30,414 unclipped subyearling Chinook salmon (19.1%), 53,798 clipped steelhead (33.8%), 11,872 unclipped steelhead (7.5%), 558 unclipped sockeye/kokanee salmon (0.4%), and 2,295 coho salmon (1.4%). There were an estimated 2,304 Chinook salmon fry (1.4%) collected during 2024. Post-season bypass occurred from October 1 through December 15.

By May 1, 51.3% of the total yearly collection for 2024 had arrived. The percent of the total collection arriving by the end of June and the end of July was 90.6% and 99.9%, respectively. Juvenile salmonids passing through the LMN JFF in August and September contributed just 0.07% of the 2024 season collection total and consisted primarily of unclipped subyearling Chinook salmon.

In 2024, the peak daily collection total and date for each species group were 6,724 clipped yearling Chinook salmon (March 29), 980 unclipped yearling Chinook salmon (April 22), 1,597 clipped subyearling Chinook salmon (July 10), 4,484 unclipped subyearling Chinook salmon (July 10), 4,325 clipped steelhead (April 24), 1,020 unclipped steelhead (May 18), 140 unclipped sockeye/kokanee salmon (May 19), and 350 coho salmon (April 30). Total daily collection in 2024 peaked at 6,992 smolts on March 29. Peak collection date and daily collection total by species group are listed in Table 4.

Table 3. Annual collection, bypass, and transport at Lower Monumental Dam, 2020-2024.

Year	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/ Unclip	
Collection										
2020	378,434	97,869	53,739	113,590	195,230	57,550	8,145	214	18,451	923,222
2021	104,155	17,085	27,414	56,418	68,200	22,481	2,042	1564	7,488	306,847
2022	667,324	119,881	103,534	131,433	366,940	131,149	17,045	1250	23,040	1,561,596
2023	984,774	186,631	45,882	106,814	585,500	174,946	15,404	1,325	12,775	2,114,051
2024	36,009	6,146	15,813	30,414	53,798	11,872	0	558	2,295	159,209 ¹
5-Yr Avg	434,139	85,522	49,276	87,734	253,934	79,600	8527.2	982.2	12,810	1,012,524
Bypass										
2020	4,030	1,877	26085	65,890	2,799	739	-	10	287	101,717
2021	20,522	1,096	17,923	41,511	13,984	1791	8	72	598	97,505
2022	18,561	9,377	71,223	92,120	23,492	10,367	295	150	7290	232,875
2023	236,774	39,374	6,181	12,482	139,984	35,635	-	1	127	470,558
2024	20,367	3,736	8,917	19,978	15,142	3,644	-	51	624	74,724 ²
5-Yr Avg	60,051	11,092	26,066	46,396	39,080	10,435	101	56.8	1,785	195,023
Truck										
2020-2024	---	---	---	---	---	---	---	---	---	---
Barge										
2020	374,298	95,939	27,619	47,643	192,349	56,792	8,142	204	18,157	821,143
2021	83,586	15,976	9,482	14,887	54,167	20,679	2,029	1490	6,885	209,181
2022	648,440	110,397	32,297	39,295	343,344	120,754	16,749	1,091	15,747	1,328,114
2023	746,835	147,019	39,632	82,183	445,224	139,261	15,398	1,323	12,741	1,629,616
2024	15,606	2,402	6,860	10,389	38,597	8,209	0	505	1,670	84,238
5-Yr Avg	373,753	74,347	23,178	38,879	214,736	69,139	8,464	923	11,040	814,458
Total Transported										
2020	374,298	95,939	27,619	47,643	192,349	56,792	8,142	204	18,157	821,143
2021	83,586	15,976	9,482	14,887	54,167	20,679	2,029	1490	6,885	209,181
2022	648,440	110,397	32,297	39,295	343,344	120,754	16,749	1,091	15,747	1,328,114
2023	746,835	147,019	39,632	82,183	445,224	139,261	15,398	1,323	12,741	1,629,616
2024	15,606	2,402	6,860	10,389	38,597	8,209	0	505	1,670	84,238
5-Yr Avg	373,753	74,347	23,178	38,879	214,736	69,139	8,464	923	11,040	814,458

Table 3. Annual peak collection dates at Lower Monumental Dam, 2020-2024.

Year	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip\Unclip	
2020	May-17	May-23	Jul-03	Jul-03	Apr-26	May-08	May-17	May-19	May-22	Apr-26
	31,200	5,300	3,100	6,660	25,300	5,400	1,600	100	1,700	46,600
2021	9-May	9-May	30-Jun	30-Jun	22-Apr	9-May	14-May	2-May	8-May	9-May
	14,050	2,050	4,482	10,415	5,257	1,950	320	220	725	21,700
2022	9-May	8-May	10-Jun	9-Jun	9-May	18-May	19-May	16-May	6-Jun	9-May
	74,400	9,400	7,200	8,800	39,600	13,600	6400	400	1650	127,200
2023	5-May	6-May	17-May	13-May	4-May	5-May	16-May	14-May	22-May	5-May
	112,800	16,800	3,203	7,800	96,525	17,600	5,400	400	2,200	223,600
2024	29-Mar	22-Apr	10-Jul	10-Jul	24-Apr	18-May	26-Mar	19-May	30-Apr	29-Mar
	6,724	980	1,597	4,484	4,325	1,020	0	140	350	6,992




Figure 2. Daily juvenile salmonid collection, all species combined, versus daily average river flow at Lower Monumental Dam, 2024.

Sampling

Sampling is defined as diverting and segregating groups of fish in a consistent fashion so data collected from those segregated groups will accurately represent all fish collected. Fish were sampled at LMN to monitor fish condition, ensure the collection system is operating correctly, and to train personnel on facility operation and sampling protocols. This type of sampling is termed “sampling for condition”, or “condition sampling”. EAS staff aim to condition sample 100 individuals of each species present in that day’s sample in accordance with the Fish Passage Center (FPC) condition sampling protocols. Individuals present in the sample that are not part of the condition sample are enumerated and examined for descaling only then routed to transport areas or bypassed, depending on which mode the JFF is operating in.

Collection for fish condition sampling began at 0700 hours on March 25 and continued until 0700 hours on October 1. Sampling occurred at 0700 until April 23, when a partial sample occurred at 1300, from a collection period of 0700 to 1300 to transition to every-day sampling on a 1300 to 1300 schedule for transport season (April 24 to June 20). Daily fish sampling concluded on June 20, at which point the barging season had ended and sampling was moved to an every-other-day schedule. Barge transport ended on 17 June but fish were sampled daily until 20 June. Collection/sampling was suspended twice due to high river temperatures during the 2024 season, on July 22 and August 20.

Total sampling includes both “condition sampling” as well as “non-condition sampling,” which was conducted during the 2024 operating year. A total of 17,559 fish (11.2% of the total collection) were sampled in 2024. Within each species group, the number and percent sampled of those collected in that group were 3,736 clipped yearling Chinook salmon (10.4%), 571 unclipped yearling Chinook salmon (9.3%), 2,601 clipped subyearling Chinook salmon (16.4%), 5,052 unclipped subyearling Chinook salmon (16.6%), 4,098 clipped steelhead (7.6%), 1,192 unclipped steelhead (10.0%), 53 unclipped sockeye/kokanee salmon (9.5%), and 256 coho salmon clipped and unclipped (11.2%) (Table 5).

Average weekly sample rates can be found in Table 6 and ranged from 0.3% to 100.00%.

Table 4. Annual percentage sampled out of total collection for each juvenile salmonid species group at Lower Monumental Dam, 2020-2024.

Year	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/Unclip	
2020	1.3	2.4	8.9	9.8	1.9	1.7	0.8	1.9	1.9	3.1
2021	4.3	9.2	13.5	11.8	4.4	5.2	4.9	5.5	7.5	6.9
2022	0.7	1.1	2.4	3.1	1	0.9	0.3	1.4	1.4	1.1
2023	0.7	1.3	5.8	4.0	0.9	0.8	0.6	1.8	2.1	1.1
2024	10.4	9.3	16.4	16.6	7.6	10.0	--	9.5	11.2	11.2

--: No clipped sockeye salmon were encountered during the 2024 season.

Table 5. Weekly average sample rates in percent and sample totals at Lower Monumental Dam, 2024.

Week Ending	Weekly Rate (%)	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Totals*
		Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/Unclip	
28-Mar	25.0	0	1	0	0	0	1	0	0	0	2
4-Apr	13.3	2,396	59	5	5	2	12	0	0	0	2,479
11-Apr	12.5	135	10	0	1	41	6	0	0	0	193
18-Apr	16.7	173	173	1	0	528	114	0	0	0	989
25-Apr	5.0	218	148	1	5	772	106	0	0	2	1,252
2-May	3.1	356	59	0	1	764	108	0	0	30	1,318
9-May	23.2	74	16	0	1	299	58	0	0	13	461
16-May	21.7	163	26	2	15	506	159	0	0	43	914
23-May	7.9	171	31	108	145	664	325	0	35	34	1,513
30-May	19.3	18	7	322	422	184	162	0	11	43	1,169
6-Jun	14.3	13	11	602	949	124	78	0	4	38	1,819
13-Jun	11.4	8	11	255	447	135	38	0	1	24	919
20-Jun	31.4	6	5	347	584	48	9	0	0	10	1,009
27-Jun	33.3	3	8	356	566	19	13	0	2	12	979
4-Jul	22.5	1	2	221	591	3	1	0	0	2	821
11-Jul	10.0	0	0	292	809	1	1	0	0	2	1,105
18-Jul	18.8	0	0	51	189	1	0	0	0	2	243
25-Jul	37.5	0	0	32	214	2	0	0	0	0	248
1-Aug	66.7	0	0	1	24	0	0	0	0	1	26

Week Ending	Weekly Rate (%)	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Totals*
		Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/ Unclip	
8-Aug	100.0	1	4	1	34	4	0	0	0	0	44
15-Aug	100.0	0	0	2	11	0	1	0	0	0	14
22-Aug	100.0	0	0	1	2	1	0	0	0	0	4
29-Aug	100.0	0	0	0	11	0	0	0	0	0	11
5-Sep	100.0	0	0	0	10	0	0	0	0	0	10
12-Sep	100.0	0	0	1	7	0	0	0	0	0	8
19-Sep	83.3	0	0	0	3	0	0	0	0	0	3
26-Sep	75.0	0	0	0	3	0	0	0	0	0	3
3-Oct	50.0	0	0	0	3	0	0	0	0	0	3
Total Sampled		3,736	571	2,601	5,052	4,098	1,192	0	53	256	17,559
% of Sample		21.3	3.3	14.8	28.8	23.3	6.8	n/a	0.3	1.5	100
% of Collection		10.4	9.3	16.4	16.6	7.6	10.0	n/a	9.5	11.2	11.2

* Daily 24-hour sampling at LMN began on April 23 this season.

-- : No sampling occurred due to warm water temperatures.

Transportation-Barge Loading Operations

An estimated 84,238 juvenile salmonids (52.9% of the collection total) were transported from LMN in 2024. All were transported by barge; no fish were trucked in 2024. Within each species group, the number transported and percent of those collected in each group was 15,606 clipped yearling Chinook salmon (43.3%), 2,402 unclipped yearling Chinook salmon (39.1%), 6,860 clipped subyearling Chinook salmon (43.4%), 10,389 unclipped subyearling Chinook salmon (34.2%), 38,597 clipped steelhead (71.7%), 8,209 unclipped steelhead (69.1%), 505 unclipped sockeye/kokanee salmon (90.5%), and 1,670 clipped and unclipped coho salmon (72.8%) (Table 3).

Fish were transported by barge from April 24 through May 4, May 18 to June 1, then every fourth day from June 1 to June 17. The fish passage plan requires fish to be transported from April 24 to June 20, but barging was halted twice in 2024 due to high flow.

Daily barge transport was halted at Lower Monumental with the May 3 barge. The barge dock was damaged during tie up on May 2. Bracing the deck I-beams between the middle and upstream coffer cells was pushed forward and no longer contacts the decking I-beams making it unsafe to load fish onto the barge by the Lower Monumental staff. Safety measures for loading the barge were evaluated and implemented and barging resumed on May 18.

On May 31 the Regional Implementation Oversight Group determined transport at Lower Monumental would occur every fourth day. Transport continued to occur on June 1 then the fish were bypassed on June 2 and 3. Collection for barging began again on June 4 and fish were barged on June 5. This four day pattern continued for the rest of the barging season, the need to stop spill at Lower Monumental Dam when the fish barge transits the tailrace causes flow fluctuations and the need for raised MOP at Ice Harbor Dam, there were indications of adult fish passage delay at Ice Harbor Dam and some regional fisheries managers believe the flow and minimum operating pool fluctuations were the cause of the passage delay.

Transportation-Truck Loading Operations

Juvenile fish were scheduled to be transported by truck from August 1 to October 1. Per the 2024 Fish Operations Plan, the Lower Monumental trucking schedule is contingent upon the Technical Management Team approval; the Technical Management Team decision was that no truck transport would take place from LMN during the 2024 transport season.

Bypass

During the 2024 season, a total of 74,724 fish were bypassed (46.9% of the collection total) (Table 3). Within each species group, the number bypassed and percent of those collected in each group was 20,367 clipped yearling Chinook salmon (56.6%), 3,736 unclipped yearling Chinook (60.8%), 8,917 clipped subyearling Chinook salmon (56.4%), 19,978 unclipped subyearling Chinook salmon (65.7%), 15,142 clipped steelhead (28.1%), 3,644 unclipped steelhead (30.7%), 51 unclipped sockeye salmon (9.1%), and 624 coho salmon (27.2%). An estimated 2,265 Chinook salmon fry were bypassed during 2024. No clipped sockeye/kokanee salmon were bypassed for the 2024 season. These numbers do not include fish bypassed by the PIT-tag diversion system.

Juvenile salmonids were bypassed rather than transported for the following purposes this season:

- Condition sampling and secondary bypass occurred (see condition sampling frequency in sampling section).
- Salmonid fry measuring less than 60 millimeters (mm) were bypassed and not sampled due to smolt monitoring protocol.
- The PTAGIS database revealed 31,945 PIT-tagged fish of different species groups were bypassed through the PIT-tag system. These fish are not included in the facility bypass total. PIT-tag diversion gates are set to bypass PIT-tagged fish when sample rates are 20% or higher and during sampling

intervals when fish are being collected for research to prevent anesthetizing study fish a second time.

The fish rearing designation used by PTAGIS is hatchery/wild, not clipped/unclipped; therefore, the hatchery/wild designation is used to report the PIT-tag numbers in the following section rather than the clipped/unclipped designation used throughout the rest of this report. According to the PTAGIS database the total of PIT-tagged fish that were detected in the JFF was 53,304. The total by unit group was as follows: Chinook salmon (17,545 hatchery and 2,593 wild), 20 steelhead (8,966 hatchery, 1,863 wild and 4 of unknown designation), 777 sockeye salmon (771 hatchery and 6 wild), 164 coho salmon (164 hatchery), and 31 fish of unknown species. An unknown number of other fish were bypassed incidentally with the PIT-tagged fish as the PIT-tag diversion gates opened and closed to divert the PIT-tagged fish.

Migration, Sampling, and Bypass of Juvenile Lamprey

Pacific lamprey, *Entosphenus tridentatus*, the primary species found at LMN and are characterized by the presence of three large teeth and posterior teeth on the oral disc. Pacific lamprey spawn in similar habitats to salmon. Spawning occurs between March and July, depending upon location within their range.

Metamorphosis from the larvae stage (ammocoetes) to the juvenile stage (macrophthalmia) occurs over a period of several months. During this time, they develop eyes, teeth, and become free swimming. They drift and swim downstream as they migrate to the ocean. It is the macrophthalmia stage when most of the lamprey end up in the sample collection at the JFF. Sample data for Pacific lamprey juvenile life stages are presented in Table 7.

Table 6. Pacific lamprey in the sample and separator, and the total collection for both life stages, with mortality 2020–2024.

Year	Sample	Separator	Mortalities	Total Collection
Pacific Lamprey (Macrophthalmia)				
2020	37,361	2	1	37,364
2021	2,930	0	1	2,931
2022	305,417	13	10	305,440
2023	509,829	13	46	509,888
2024	1,394	1,309	98	13,791
Pacific Lamprey (Ammocoete)				
2020	388	1	0	389
2021	1096	0	0	1,096
2022	20,041	4	18	20,063
2023	4,159	2	5	4,166
2024	9	9	1	137

Incidental Species

Non-target fish species that were too large to pass through the separator bars were recorded and bypassed through the adult release pipe at the separator. Those small enough to pass through the separator bars were either sampled and bypassed, or held in the raceways. Raceway incidentals are bypassed to the river. Fortunately, most incidental fish generally arrive late in the season when a high percentage of the collection is sampled. At this time, incidental species are removed while working up the sample, therefore avoiding transport. Sample fish from each incidental species were counted and their total numbers were calculated using the sample rate. These numbers were then added with separator counts of the same group to estimate the total collection for each species. The most common incidental species groups for 2024 included the following: juvenile American shad, *Alosa sapidissima* (38,343); juvenile Pacific lamprey macrophthalmia (13,791); Siberian prawn, *Exopalaemon modestus* (2,921); walleye, *Sander vitreus* (539); sculpin, *Cottus* spp. (294); rainbow trout, *Oncorhynchus mykiss* (158); and larval Pacific lamprey ammocoete (137).

In general, incidental fish collection numbers were much lower during the 2024 operating season compared to the 2023 season. The number of juvenile shad, 38,343, in 2024 is lower than the 47,759 collected in the 2023 operating year. In the 2024 season, 13,791 juvenile Pacific lamprey macrophthalmia were collected, compared to 509,842 in the 2022 season. Pacific lamprey ammocoete numbers decreased from 4,161 in 2023 to 137 in 2024. Walleye numbers were also reduced compared to 2023 collection numbers. Approximately 539 walleye were collected in the 2024 season, while 1,221 were collected in the 2023 season. Estimated numbers of some groups may also become exaggerated high or low due to the low sample rates at the time of their collection. A summary of incidental fish collection during 2023 sampling efforts is provided in Table 8.

Table 7. Estimated collection of incidental species at Lower Monumental Dam, 2024.

Common Name	Scientific Name	Sample	Separator	Total 2024 Collection ^a
American shad (Adult)	<i>Alosa sapidissima</i>	3	62	66
American shad (Juvenile)	<i>A. sapidissima</i>	26,810	9,970	38,343
Bluegill and Pumpkinseed	<i>L. macrochirus/ gibbosus</i>	12	5	51
Bullhead (misc.)	<i>Ameiurus spp.</i>	2		5
Bull Trout	<i>Salvelinus confluentus</i>			0
Channel catfish	<i>Ictalurus punctatus</i>	19		24
Chiselmouth	<i>Acrocheilus alutaceus</i>			0
Common carp	<i>Cyprinus carpio</i>	2	5	7
Crappie	<i>Pomoxis spp.</i>	24	3	37
Kokanee	<i>Oncorhynchus nerka</i>			0
Northern Pikeminnow	<i>Ptychocheilus oregonensis</i>		1	1
Pacific lamprey (Adult)	<i>Lampetra tridentatus</i>	34	17	70
Pacific lamprey (Juvenile)	<i>L. tridentatus</i>	1,394	1,309	13,791
Pacific lamprey (Ammocoete)	<i>L. tridentatus</i>	9	9	137
Peamouth	<i>Mylocheilus caurinus</i>	1		2
O. mykiss other (Rainbow Trout)	<i>Oncorhynchus mykiss</i>	45	12	158
Sandroller	<i>Percopsis transmontana</i>			0
Sculpin	<i>Cottus spp.</i>	61		294
Siberian Shrimp/Prawn	<i>Exopalaemon modestus</i>	1,486	32	2,921
Largemouth/Smallmouth bass	<i>Micropterus dolomieu/salmoides</i>	35	10	72
Sucker (misc.)	<i>Catostomus spp.</i>	12	4	61
Whitefish	<i>Prosopium spp.</i>	1		4
White Sturgeon	<i>Acipenser transmontanus</i>		3	3
Walleye	<i>Sander vitreus</i>	198	56	539
Warmouth	<i>L. gulosus</i>	1		25
Yellow perch	<i>Perca flavescens</i>	15	14	108
Others			2	2
Total		30,164	11,514	56,721

^a Incidental species collection estimates are based on sampled number of group expanded by the sample rate plus separator count.

Adult Fallbacks

A total of 193 adult salmonids fell back through the juvenile bypass system and were bypassed from the separator between March 26 and October 1. The total includes 44 adult Chinook salmon, 13 jack Chinook salmon, 46 clipped steelhead, 47 unclipped steelhead, 30 sockeye salmon, and 13 coho salmon (Table 9).

The most common adult salmonid fallbacks in the 2024 season were steelhead (48.2%) with the month of September having the most steelhead fallbacks (Table 10). Total monthly adult fallbacks for all salmonid species also peaked in September, with 42.5% of all fallbacks being observed during the month of May.

Table 8. Annual totals of adult salmonids released from the separator at Lower Monumental Dam, 2020–2024.

Year	Chinook	Chinook Jack	Steelhead Clipped	Steelhead Unclip	Sockeye	Coho	Total
2020	83	44	79	112	21	8	347
2021	46	20	23	49	5	5	148
2022	125	23	153	114	2	1	418
2023	87	29	135	248	8	3	510
2024	44	13	46	47	30	13	193

Table 9. Monthly totals of adult salmonids released from the separator at Lower Monumental Dam, 2024.

Month	Chinook	Chinook Jack	Steelhead Clipped	Steelhead Unclip	Sockeye	Coho	Total
March	0	0	0	0	0	0	0
April	0	0	8	6	0	0	8
May	12	1	2	11	0	0	21
June	11	1	7	14	1	0	34
July	10	0	2	3	17	0	28
August	4	0	4	5	8	1	23
September	7	10	28	21	4	12	82
Total	44	13	46	47	30	13	193

The condition of adult salmonids was evaluated as the fish were released from the separator. Their condition was predominantly good to fair with 97.9% of the fallbacks rated in these categories (Table 11). Condition ratings of the adults examined were as follows: 177 good (91.7%), 12 fair (6.2%), 3 poor (1.6%), and 1 dead (0.5%). The one fallback mortality during the 2024 season was an unclipped steelhead.

Table 10. Condition of adult salmonids released from the separator at Lower Monumental Dam, 2024.

Condition	Chinook	Chinook Jack	Steelhead	Steelhead	Sockeye	Coho	Total
			Clipped	Unclip			
Good	39	13	44	42	26	13	177
Fair	3	0	1	4	4	0	12
Poor	2	0	1	0	0	0	3
Dead	0	0	0	1	0	0	1
Total	44	13	46	47	30	13	193

Separator Efficiency

The separator is designed with bar spacing to allow only smaller smolts—subyearling Chinook and sockeye salmon—to divert to the A-side of the collection facility. Larger smolts—steelhead and yearling Chinook salmon—divert to the B-side through wider spaced bars. Separator efficiency for 2024 by species group

was as follows: clipped yearling Chinook salmon (54.3%), unclipped yearling Chinook salmon (40.2%), subyearling Chinook salmon (53.9%), clipped steelhead (80.8%), unclipped steelhead (74.3%), and unclipped sockeye/kokanee salmon (28.3%) (Table 12). Due to staffing issues The Idaho Department of Fish and Game did not clip sockeye for the 2024 season, no clipped sockeye salmon were observed at the juvenile fish facility at Lower Monumental Dam.

Table 11. Annual separator efficiency in percent at Lower Monumental Dam, 2020–2024.

Year	Yearling Chinook		Subyearling Chinook	Steelhead		Sockeye	Sockeye
	Clipped	Unclip	Clipped/Unclip	Clipped	Unclip	Clipped	Unclip
	A-side	A-side	A-side	B-side	B-side	A-side	A-side
2020	71.9	65.1	42.4	44.8	26.9	34.5	56.4
2021	67.6	55.3	29.2	69.0	66.8	60.4	29.9
2022	71.4	57.9	50.1	49.9	76.9	54.6	49.8
2023	57.3	51.6	49.5	55.2	89.2	73.9	41.2
2024	54.3	40.2	53.9	80.8	74.3	--	28.3

Fish Condition

Descaling

Descaling data were collected from all live sample fish (full sample) rather than just a portion (subsample). Full-sample data collection provides a larger sample size and therefore a better representation of fish condition.

The descaling rate for all fish sampled in 2024 was 0.7%. The annual descaling rate by species group were as follows: clipped yearling Chinook salmon (0.6%), unclipped yearling Chinook salmon (0.8%), clipped subyearling Chinook salmon (0.7%), unclipped subyearling Chinook salmon (0.6%), clipped steelhead (0.8%), unclipped steelhead (1.1%), unclipped sockeye/kokanee salmon (1.9%), and coho salmon (0.8%) (Table 13). No clipped sockeye salmon were collected for 2024.

In 2024, the highest weekly descaling rate for all species combined was 25.0% for the week ending

August 22 (with 4 fish sampled in a week of condition sampling), while weekly descaling (0.0%) occurred 32% of weeks during the sampling season (Table 14).



Table 13. Annual descaling rates in percent for fish sampled at Lower Monumental Dam, 2020–2024.

Year	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/Unclip	
2020	1.5	1.6	1.6	1.1	2.5	2.9	2.4	0	2.9	1.9
2021	1.0	1.6	1.2	1.4	3.2	3	4.1	4.7	3.6	1.7
2022	1.2	1.6	1.4	2.2	2.2	1.7	0.0	0.0	2.3	1.7
2023	0.4	0.5	0.1	0.4	1.9	3.6	0.0	0.0	0.4	0.9
2024	0.6	0.8	0.7	0.6	0.8	1.1	--	1.9	0.8	0.7
Average	0.9	1.2	1.0	1.1	2.1	2.5	1.6	1.3	2.0	1.4

Table 12. Weekly descaling rates in percent for fish sample at Lower Monumental Dam, 2024.

Week Ending	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Both	
	CCH1	UCH1	CCH0	UCH0	CST	UST	CSO	USO	CO	
28-Mar	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
4-Apr	0.21	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.20
11-Apr	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
18-Apr	2.65	2.45	0.00	0.00	1.37	0.89	0.00	0.00	0.00	1.71
25-Apr	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2-May	1.72	0.00	0.00	0.00	0.44	0.99	0.00	0.00	0.00	0.83
9-May	0.00	0.00	0.00	0.00	0.46	0.00	0.00	0.00	0.00	0.29
16-May	0.68	0.00	0.00	0.00	0.22	0.74	0.00	0.00	0.00	0.37
23-May	1.19	0.00	0.00	0.00	0.97	1.68	0.00	0.00	2.94	0.99
30-May	0.00	0.00	0.32	0.00	1.92	1.44	0.00	9.09	2.33	0.74
6-Jun	0.00	0.00	0.53	0.34	0.00	1.39	0.00	0.00	0.00	0.41
13-Jun	12.50	0.00	0.81	0.23	4.84	3.03	0.00	0.00	0.00	1.25
20-Jun	0.00	0.00	0.59	0.35	4.26	0.00	0.00	0.00	0.00	0.60
27-Jun	0.00	0.00	0.31	0.73	0.00	0.00	0.00	0.00	0.00	0.54
4-Jul	0.00	0.00	1.06	1.55	0.00	0.00	0.00	0.00	0.00	1.40

Week Ending	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Both	
	CCH1	UCH1	CCH0	UCH0	CST	UST	CSO	USO	CO	
11-Jul	0.00	0.00	1.97	0.68	0.00	0.00	0.00	0.00	0.00	1.00
18-Jul	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
25-Jul	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1-Aug	0.00	0.00	0.00	5.00	0.00	0.00	0.00	0.00	0.00	4.55
8-Aug	100.00	0.00	0.00	2.94	0.00	0.00	0.00	0.00	0.00	4.55
15-Aug	0.00	0.00	50.00	9.09	0.00	0.00	0.00	0.00	0.00	15.38
22-Aug	0.00	0.00	0.00	0.00	100.00	0.00	0.00	0.00	0.00	25.00
29-Aug	0.00	0.00	0.00	9.09	0.00	0.00	0.00	0.00	0.00	9.09
5-Sep	0.00	0.00	0.00	10.00	0.00	0.00	0.00	0.00	0.00	10.00
12-Sep	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
19-Sep	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
26-Sep	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
3-Oct	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Total Descaled	20	4	17	28	30	12	---	1	2	114
Total Examined	3,619	529	2,417	4,734	3,684	1,070	---	52	255	16,360
% Descaled	0.6	0.8	0.7	0.6	0.8	1.1	---	1.9	0.8	0.7

Other Injury and Disease

Injury and disease data were collected from a subsample goal of 100 of the dominant species and not more than 100 each of the non-dominant species. A total of 9,662 fish were examined for condition. The most common symptoms observed in 2024 were fin injury (1,316) and fin hemorrhage (425).

Blood pooling is defined as the vasodilatation of the capillaries in fins (also referred to as pink fin). It seems to be a symptom of anesthetic use during higher water temperatures and is mostly found on subyearling Chinook salmon. Evidence of blood pooling was found on 0.45% of all fish examined. The incidence of blood pooling by species group was as follows: yearling Chinook salmon 0.16%, subyearling Chinook salmon 0.51%, steelhead 0.53%, sockeye/kokanee salmon 1.92%. No blood pooling was observed on coho salmon for the 2024 season.

Fin injuries were found on 13.62% of all fish examined. The incidence of fin injury was as follows: yearling Chinook salmon 34.05%, subyearling Chinook salmon 5.33%, steelhead 13.37%, and sockeye/kokanee salmon 9.62%, and coho salmon 4.71%. Fin hemorrhaging often coincided with split fin injuries.

Fin hemorrhaging is the discharge of blood outside the fin tissue. Fin hemorrhaging is a sign of trauma and was found on 4.40% of all fish examined for injuries. The incidence of fin hemorrhaging by species group was as follows: yearling Chinook salmon 5.65%, subyearling Chinook salmon 7.02%, and steelhead 0.38%, sockeye/kokanee salmon 3.85%, and coho salmon 1.57%.

Bird marks were observed on 0.75% of all fish examined. The incidence of bird marks was as follows: yearling Chinook salmon 0.90%, subyearling Chinook salmon 0.47%, steelhead 1.04%, sockeye/kokanee salmon 1.92%, and coho salmon 0.39%. Predatory fish marks were found on 0.92% of all fish examined. The incidence of fish marks was as follows: yearling Chinook salmon 1.37%, subyearling Chinook salmon 0.87%, and steelhead 0.72% sockeye/kokanee salmon 3.58%, and coho salmon 0.39%. Lamprey marks were seen on 0.52% of all fish examined. The incidence of lamprey marks were as follows: yearling Chinook salmon 0.05%, subyearling Chinook salmon 0.56%, and steelhead 0.72% sockeye/kokanee salmon 1.92%, and coho salmon 0.39%.

Fungus was found on 0.65% of all fish examined. Fungus on fish was often found concurrently with body injuries. The incidence of fungus was as follows: yearling Chinook salmon 0.37%, subyearling Chinook salmon 0.02%, steelhead 1.69%, and sockeye/kokanee salmon 1.92%. No fungus was observed in 2024 for coho salmon.

Columnaris was not observed 2024. It occurs most frequently in subyearling Chinook salmon but has been seen on coho salmon and steelhead as well. Typically, it is found on the fish during the warmer water conditions of July, August, and September. Columnaris can be recognized by the presence of yellowish lesions on the belly, as well as some damage to the gills, pelvic fins, snout, and caudal fins. It has also been found in the dorsal region.

Mortality

Annual facility mortality for all groups combined was 0.10% in 2024 (Table 15) and totaled 158 fish. Within each species group, the number of facility mortalities and percent of those collected in that group was 20 clipped yearling Chinook salmon (0.06%), 6 unclipped yearling Chinook salmon (0.10%), 14 clipped subyearling Chinook salmon (0.09%), 25 unclipped subyearling Chinook salmon (0.08%), 54 clipped steelhead (0.10%), 11 unclipped steelhead (0.09%), 1 unclipped sockeye/kokanee salmon (0.18%), and no coho salmon (0.0%). No clipped sockeye/kokanee salmon were collected in 2024, and there were 27 chinook fry facility mortalities. In 2024, the highest rate of mortality was 0.22% on the week ending Jun 7 and the lowest rate of mortality was 0%, occurring on multiple weeks (Table 16).

Annual sample mortality for all groups combined in 2024 was 0.50% and totaled 88 fish. The number of sample mortalities and mortality rate by species group was 16 clipped yearling Chinook salmon (0.43%), 2 unclipped yearling Chinook salmon (0.35%), 22 clipped subyearling Chinook salmon (0.85%), 22 unclipped

subyearling Chinook salmon (0.44%), 5 clipped steelhead (0.12%), 8 unclipped steelhead (0.67%), 1 unclipped sockeye kokanee (1.89%), and no coho salmon (0.39%) (Table 17). There were 12 chinook fry sample mortalities (4.7%).

Table 13. Annual facility mortality in percent at Lower Monumental Dam, 2020-2024.

Year	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/Unclip	
2020	0.03%	<0.01%	0.07%	0.05%	0.04%	0.03%	0.04%	--	0.04%	0.03%
2021	0.05%	0.08%	0.03%	0.04%	0.07%	0.05%	0.24%	<0.01%	0.07%	0.05%
2022	0.05%	0.09%	0.01%	0.01%	0.03%	0.02%	0.01%	0.80%	0.01%	0.04%
2023	0.11%	0.12%	0.15%	0.05%	0.04%	0.03%	0.04%	0.08%	0.02%	0.08%
2024	0.06%	0.10%	0.09%	0.08%	0.10%	0.09%	--	0.18%	0.00%	0.10%

Table 16. Weekly facility mortality in percent at Lower Monumental Dam, 2024.

Week Ending	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/Unclip	
	CCH1	UCH1	CCH0	UCH0	CST	UST	CSO	USO	CO	
28-Mar	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
4-Apr	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
11-Apr	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
18-Apr	0.3	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.11
25-Apr	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.00
2-May	0.1	0.3	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.05
9-May	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.00
16-May	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.00
23-May	0.2	0.0	0.2	0.2	0.0	0.1	0.0	0.0	0.0	0.08
30-May	0.8	0.0	0.2	0.1	0.0	0.1	0.0	1.7	0.0	0.12
6-Jun	0.0	0.0	0.2	0.3	0.0	0.2	0.0	0.0	0.0	0.21
13-Jun	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.03
20-Jun	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.00
27-Jun	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.00
4-Jul	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.06
11-Jul	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
18-Jul	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
25-Jul	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1-Aug	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Week Ending	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/Unclip	
	CCH1	UCH1	CCH0	UCH0	CST	UST	CSO	USO	CO	
8-Aug	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15-Aug	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
22-Aug	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
29-Aug	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5-Sep	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
12-Sep	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
19-Sep	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
26-Sep	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
3-Oct	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Table 14. Annual sample mortality in percent at Lower Monumental Dam, 2020-2024.

Year	Yearling Chinook		Subyearling Chinook		Steelhead		Sockeye		Coho	Total
	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clipped	Unclip	Clip/Unclip	
2020	0.20%	0.30%	0.30%	0.40%	0.20%	0.30%	1.50%	0.00%	0.00%	0.30%
2021	0.60%	0.60%	0.20%	0.20%	0.40%	0.40%	2.00%	1.20%	0.40%	0.70%
2022	0.70%	0.80%	0.30%	0.20%	0.30%	0.40%	0.00%	0.00%	0.00%	0.40%
2023	0.50%	0.10%	0.50%	0.40%	0.70%	0.40%	2.30%	0.00%	0.80%	0.50%
2024	0.43%	0.35%	0.85%	0.44%	0.12%	0.67%	---	1.89%	0.39%	0.50%

Research

Gas Bubble Trauma

Juvenile Chinook salmon and steelhead were sampled once a week for GBT from April 7 to July 31. This season, 1,124 fish were sampled for GBT. ODFW personnel examined up to 102 individuals of each of the following groups: yearling Chinook salmon, subyearling Chinook salmon, and juvenile steelhead. The fish were examined for evidence of bubbles in paired and unpaired fins and in the eye, as per FPC GBT protocols. GBT fish were bypassed to the river after examination. Weekly GBT sampling continued for up to 4 hours or until 102 fish had been sampled per species group. The number of fish sampled for GBT, by species group, was 109 clipped yearling Chinook salmon, 33 unclipped yearling Chinook salmon, 149 clipped subyearling Chinook salmon, 309 unclipped subyearling Chinook salmon, 410 clipped steelhead, and 114 unclipped steelhead. In the 2024 season, 10 fish showed signs of GBT in the fins (0.88%).

PNNL Lamprey Study

Pacific Northwest National Laboratory (PNNL) has been contracted by the U.S. Army Corps of Engineers (USACE) – Walla Walla District to perform a dam passage and survival study of juvenile Pacific lamprey through Federal Columbia River Power System hydroelectric projects on the Snake River. The study will evaluate lamprey passage and survival and performance of dam configuration and operations from the forebay of Lower Granite Dam (LWG) on the Snake River downstream to the mouth of the Snake River near its confluence with the Columbia River. The study seeks to implement a plan to continue to improve Pacific lamprey passage conditions at USACE dams. PNNL used Juvenile Salmon Acoustic Telemetry System (JSATS) Eel and Lamprey Acoustic Tags and nodes in the river to track the fish. Fish are released 20 kilometers above LWG at Blyton Landing.

Steelhead Kelt Study

Efforts to recover declining steelhead stocks within the Columbia River Basin have ranged from harvest reduction, habitat restoration, passage improvements at mainstem Columbia River hydropower facilities, and hatchery propagation. More recently, the use of kelt reconditioning has been investigated to increase total reproductive potential of steelhead populations (Hatch et al. 2020). Kelt reconditioning consists of the collection of post-spawned steelhead and the administration of prophylactics and feed for the purpose of improving survival relative to the untreated condition. Upon release, these fish are intended to return to natal populations, thereby increasing escapement and productivity if reconditioned individuals successfully spawn. Collection for the Nez Perce steelhead kelt study and rehabilitation began in early April, once set up of the collection tank was completed. On July 20, all the study equipment was removed for the remainder of the year, except for the holding tank, which will remain until the 2023 collection year when the study begins again.

Facility Operations and Maintenance

Turbine Operations

Efforts were made to operate all turbine units within 1% of peak efficiency from April 3 to October 31. Deviations were infrequent and brief or required by the Bonneville Power Administration. Unit priority was in effect from March 1 to November 30. Units were taken OOS for various reasons throughout the year. Table 18 below provides a summary of unit outages and causes.

Table 15. Unit outages and causes at Lower Monumental, 2023.

Dates out of service (OOS)	Unit	Reason out of service (OOS)
March 2	Unit 2	Faulty turbine bearing low oil flow alarm
March 3 – 8	Unit 2	Thrust Bearing pressure switch
March 15–17	All Units	Trash rack raking (6-8 hours/day alternating units)
March 21–23	All Units	STS installation
March 28	Unit 2	2B STS failure
June 6	Unit 5	5B STS OOS
June 13 – July 28	Unit 5	6-year maintenance overhaul/T2 repair
June 19 - 21	Unit 3	Annunciator breaker tripped
June 23	Unit 1	Bad Nexus board
July 19 -20	Units 1 – 4	Trash rack raking
August 1 – 18	Unit 2	Annual/Exciter Replacement
August 22	Unit 1 – 4	Preparation for T2 repair
August 22 – November 4	Unit 5	T2 repairs
August 22 – November 7	Unit 6	Annual maintenance/T2 repairs
September 26 – October 13	Unit 4	Annual maintenance
March-December monthly (2-3 days)	All Units	STS/VBS inspection/hub tapping on fixed blade units

Spill Operations

Prior to spill season, spill only occurred if it was needed for spill excess of powerhouse capacity. Limited spill through the RSW for adult steelhead passage occurred from March 1 to 31 and again from September 1 through November 15. The spring spill program began at 0001 hours April 3 and ended 0000 hours on June 20. The summer spill program began at 0001 hours on June 21 and ended at 0000 hours on August 31.

Spillways 1, 2, 6 and 7 were taken OOS at 1800 hours on April 22 for trunnion stress issues and resolution. This also caused the RSW to be removed from service at varied times for the repair work. Spillway 7 was RTS on May 4 at 0925 hours. The application of the grease line was completed on spillway 6 on May 3 at 0900 hours and then the spillway was RTS on May 4 at 0925 hours. Spillway 3 was OOS from 0700 hours to 1700 hours on May 5 while work was being done on spillway 2. Spillway 1 RTS on May 12 at 1430 hours. Spillway 2 RTS on May 10 at 1610 hours.

Spillway 7 was forced OOS at 1520 hours on May 12 for a motor coupler issue and was RTS on May 12 at 1536 hours. Spillway 5 was placed OOS on June 7 at 1212 hours due to motor coupling issues. The spillway was RTS on June 8 at 0835 hours.

During the spill seasons, spill adjustments for navigation were done as required. After spill season ended, spill only occurred if needed for excess of powerhouse capacity.

Removable Spillway Weir

The RSW went into service at 0001 hours on April 3 with the start of the spring spill program. Due to high river temperatures correlated with low river flows, the RSW was closed on August 4, except for the fall spill to assist with steelhead passage.

The RSW was taken OOS during 0630 hours to 1700 hours from April 25 to April 28 when the powerhouse crews were working on spillway 7.

Forebay Debris

Forebay debris was moderately light for the year. The maximum amount of debris observed during ladder inspections was around 3,100 yards squared, observed in the month of April.

Trash Racks

During the winter, trash rack measurements were not taken. Before the juvenile passage season, trash racks were cleaned at all slots from March 15 to 17. Approximately 35 cubic yards of debris, mostly consisting of tumbleweeds, were removed. Only one fish was observed during the trash rack cleaning, one decomposed adult shad. The trash racks were cleaned again at mid-season, July 19 to 20. Approximately 30 yards of debris was removed from the trash racks mostly consisting of logs and sticks. The majority of the debris was removed from units 1 and 2. No problems were observed during any cleanings.

Gatewells

During the season, gatewell slots were checked during ladder inspections, approximately three times per week. The gatewell drawdown benchmark measurements were taken for all units on March 24.

A small amount of residual oil/film was found in the 6A slot on April 19. An absorption pad was placed into the slot and the control room was notified. A large chunk of plastic bucket was retrieved from the 6A gatewell slot on August 8, because its size could cause an orifice plug.

In addition, during the season small amounts of woody material were noted in the gatewell slots, and the debris coverage did not exceed 50% during any inspections. No large accumulations of woody material were noted.

Submersible Traveling Screens

During the winter maintenance season, electrical cables, gearboxes, motors, and screens were examined and repaired or replaced as required. The STSs were inspected and tested on March 16 prior to installation. STSs for units 5 and 6 were deployed on March 21. Units 3 and 4 were deployed on March 22, and units 1 and 2 were deployed on March 23.

STSs are usually operated in cycle mode when the average fork length of subyearling Chinook salmon and/or sockeye salmon is greater than 120 mm, and in continuous-run mode when either is less than 120 mm. The STSs were placed in cycle-run mode when first deployed on March 18; they were changed to continuous-run mode on June 6 due to average subyearling Chinook and sockeye lengths being less than 120 mm. They were changed back to cycle-run mode on July 19 when the average lengths of collected fish were greater than 120 mm.

STS deployed in slot 2B had a motor failure on March 28 and was immediately replaced with a spare screen before the unit RTS. STS deployed in slot 3B had a motor trip with failure to reset on October 28, the unit remained OOS until November 4 after a Bonneville Power Administration (BPA) line outage for T2 repair. STS deployed in slot 5B had a motor failure on June 6 and was immediately replaced with a spare screen before the unit RTS.

The STS in the 3B slot was found not operational on October 28 at 1550 hours. Unit 3 was kept OOS due to the failure until it could be examined by powerhouse staff. Powerhouse electricians worked on the STS operating system on October 31 and November 1. After troubleshooting the STS issue, the electrical issue with the STS was fixed and the STS was returned to service on November 2.

All STSs were raised by December 15. Unit 1 had been raised prior because the unit was OOS. After the screens were raised, they were visually inspected. The STSs appeared clean and had no apparent damage to them.

Vertical Barrier Screens

VBS differentials are not measured daily at Lower Monumental due to the general structure of the VBSs. Instead, they are spot checked during the monthly STS inspections. A thorough VBS inspection occurred this year occurred July 7-8 and September 6-7. A few issues were found during the camera inspection of the VBSs in slots 4A, 4B, 5A, and 5B.

During July 11 to 12, repairs were made to the VBSs in slots 5A and 5B. Both slots required a small strip of sheet metal to be welded to the top of the VBS to diminish the gap between the gatewell concrete wall and the top of the VBS.

During the inspections on September 6 to 7, the VBS in the 4A slot appeared to be missing a sheet of metal which butts up against the concrete wall of the gatewell. The slot was dewatered on September 28, and it was discovered that the VBS was manufactured differently and did not have this additional metal sheet. There was still an obvious $\frac{1}{2}$ " gap above the VBS which was corrected by the powerhouse mechanics by fabricating a piece of sheet metal onto the VBS to prevent a future problems.

During camera inspections on September 6 to 7, the VBS in the 4B slot appeared to have a gap on the top which sticks were getting lodged. The 4B slot was dewatered and a physical inspection of the VBS occurred on September 28. After the inspection, it was determined no work was needed to the VBS. The small sticks were removed.

No other issues were found throughout the year.

Juvenile Collection Channel Orifices

During the 2023 season, the number of open orifices varied from 17 to 20 according to forebay level. The orifices were opened on March 23, after the winter maintenance period. With the Lower Monumental reservoir at minimum operating pool, water discharge through an orifice is reduced. During this period, extra orifices were opened to supply additional water to the adult fishway. Orifices were cycled and backflushed with air daily to remove debris. Orifice lights were also checked daily. If a light was not working, flow was directed to the other orifice in the slot until repairs could be made. The orifices were closed for the winter maintenance period on December 15 at 1055 hours.

The orifices at 5C29 and 6B33 in the channel were inspected after the powerhouse staff reported they appeared to be binding up. The JFF mechanics checked the cylinders and valves for any broken components on October 15 and reported them back in service to the control room.

Primary Dewatering Structure

During the winter maintenance period, the leak at the joint between the JCC and PDS that was discovered during the 2021 season, was replaced with a new rubber gasket. In addition, the sprocket, chain and cable from the screen cleaning brush were replaced.

The PDS operated from March 23 to December 15. The compressed air screen cleaner functioned well throughout the 2023 season. The PDS mechanical screen cleaner brush functioned well except for the issues listed below.

The air bubbler in zone 3 went OOS on April 1. JFF maintenance discovered a bad air flow solenoid. Powerhouse electrician replaced the solenoid and zone 3 went back into service at 0900 hours on June 23. High-water alarms went off on July 3, 6, 7, and 8 for the PDS. However, when examining the water level in the PDS, it appeared normal. It was discovered that zone 3 had too much air flow and caused the high-water sensor to trigger. The air valve was adjusted and corrected the issue.

After multiple high-water alarms received in the control room on August 17, the PDS weirs were lowered to bring the water level 0.2-0.4 ft lower to aid with the alarms in the control room. The electricians made a few

modifications to the set points on August 18. The PDS weirs were raised to a normal setting on August 19 at 0700 hours.

During the evening of September 11, the PDS screen cleaning brush stopped in the middle of the cycle. The brush was returned to park position and reset. The brush functioned effectively during the next cleaning cycles.

The air bubbler timer was increased to run every 10 minutes and the PDS screen brush timer was decreased to run only twice a week on October 15, due to the decrease in debris coming through the system. For the remainder of the season, the brush was only run manually on Mondays and Thursdays to prevent wear and tear on the brush components.

The brush timer on the brush at the PDS was modified by powerhouse electricians on December 13 to allow the brush to be set to either a 1, 3, 6 or 12 hour run time option. This decreased the over-usage on the brush. On December 15 the brush was turned off and tagged out for the winter maintenance season once the JCC was dewatered.

JUVENILE FISH FACILITY

Separator

Sudden water level drops at the separator were not a problem this year but continued to occur. Water level remained consistent at the separator with manual operation of the automated weirs of the primary dewaterer. As has been the case for the last few years, the separator was operated at a higher water level to assure no problem with exposed separator bars would occur.

PIT-Tag System

The PIT-tag system functioned well this season. There were no problems to report with the PIT-tag system.

The PIT-tag system detected 31,945 PIT-tagged fish at the JFF from April 1 to October 1. None of these PIT-tagged fish are included in the bypass numbers. Juvenile hatchery Chinook salmon, hatchery coho salmon, and hatchery steelhead in the Snake River Basin are normally designated by fin clips, usually the adipose fin, but occasionally one of the pectoral or ventral fins.

Sample

The JFF was watered up for testing at 0820 hours on March 23. The JFF and wet lab was fully watered up on March 29. Every-third-day fish condition sampling began at 0700 on April 1. The first sample was processed on April 2, and continued every third day up until April 14, after which condition sampling took place every other day. Early-season condition monitoring consisted of a 24-hour sample on target days from 0700 hours to 0700 hours. Every-other-day sampling continued until the start of transportation operations on April 23, at which time sampling was conducted every day. During this period, fish were examined and returned to the river. Every-day sampling continued up until June 9, when the fish facility switched to every-other-day sampling in accordance with the newly updated FPC high water-temperature guidelines due to Snake River water temperatures exceeding 68.5°F.

Collection for barge transport began at 0700 hours on April 23 and ended at 1300 hours on June 9. Due to high river levels and flow, June 7 fish transport was canceled at Lower Monumental, and those fish collected were released back to the river. Beginning June 10, all fish collected were sampled for condition and then bypassed. This operation continued until October 1, when the facility was returned to primary bypass. The JFF was dewatered for winter maintenance on October 3.

The sample screen which controls the switch gates stopped working on August 10 at 2030 hours. The technician on duty checked the timing of the gates with a stopwatch to ensure the gates were still collecting at an accurate percentage. The screen was able to be rebooted on August 11 at 0745 hours.

The B-side count tank pipe was found plugged on August 11 at 0615 hours. The lead biological technician and mechanics were able to unplug the pipe at 0830 hours. The plug consisted of two dead incidental fish, a walleye and a bass, and some smaller sticks.

Other Facility Issues

Nuisance bird deterrent were place up around all the PIT-tag boxes in late June to prevent the starlings from continuing to nest behind the boxes.

The JFF mechanical shop air conditioner was replaced on July 25. During the last few weeks in July, the raceway flow coming from the diffusers appeared to be less than previous months. The raceway water was lowered, and the diffuser plates were cleaned on July 25. This corrected the issue with a lower water pressure.

A power outage occurred at the fish facility on August 18 at 1615 hours that lasted approximately 5 minutes. The only noticeable effect was the outfall cannon had to be restarted.

An air cylinder for the lamprey bypass system started leaking on August 20. Air supply was turned off until the repair was made on August 31. This did not affect operations since the lamprey bypass is not used when fish are not going to the raceways.

A power outage occurred at the JFF on August 22 at 0655 and then again at 1315 to assist with the XJ breaker and T-2 rehabilitation. No issues occurred from the power outages, except the bird cannon had to be reset after each power outage. Two additional power outages occurred on August 23, with no issues caused from the outages.

The B-side PIT tank valve would not close using the electrical switch on September 12. The powerhouse electricians examined the PIT tank system that same day. The electricians believed the corrosion on the switch contacts caused the PIT tank to not function. In the afternoon, after the inspection, the B-side PIT tank was RTS.

The PIT-tag slide gate pins were replaced on both the A- and B-side on October 6.

The sample gate screen in the separator booth stopped functioning in the evening of September 12. The screen was reset twice and did not reboot correctly. The fish biologist rebooted the screen by removing the breaker and allowing it to cool for approximately 10 minutes. Once rebooted, the screen was able to be used. As a precaution since this had occurred already in the season, Pacific States Marine Fisheries Commission was notified of the issue.

A brief power outage occurred on October 31 from 0815 to 0840 hours while during the preparation for the BPA power grid modification. A loss of power occurred to the station service on November 3 from 1643 to 1830 hours. The project was not running any units during the power outage, as BPA was performing power grid modifications. The single project operator was able to restore power for approximately 30 minutes during that time while troubleshooting the power loss issue. All systems returned to service at 1830 hours once station service was able to be restored.

During the JCC dewater on December 15, the winch that lifts the gate covering access to the return to river pipe was damaged. A new winch was ordered and will be replaced during the 2023 winter maintenance period.

FISH SALVAGE

Only juvenile fish salvage at various locations will appear in this report. Fish were not always examined for clips as their survival was a higher priority.

The fish rescue occurred for the North adult fish ladder and powerhouse collection channel on January 11 at 0700 hours. There was a total of 27 live clipped adult steelhead and 5 lived adult unclipped steelhead returned to the river. Most of the steelhead were less than 20 inches in length. In addition, there were 55 adult catfish, 3 smallmouth bass, 1 walleye, 1 carp and 1 perch returned to the river at the tailrace.

Unit 4 scrollcase fish rescue took place on July 22 at 1300 hours. One catfish was released and there were no mortalities found.

A fish salvage took place from the gatewell slot of 4A. The fish salvage occurred while the maintenance bulkhead was being removed from the 4A slot. The mechanical crew realized the fish were trapped on top of the bulkhead and immediately stopped all operations. They then called the fish team who came and removed the trapped fish who were returned to the river unharmed. There were three clipped steelhead rescued during the salvage. A fish salvage occurred at unit 6 scrollcase once it was dewatered on October 6 at 0825 hours. There were no fish observed in the scrollcase.

The JCC was dewatered on December 15. The orifices were closed at 1055 hours and the channel dewatering was completed by 1230 hours. Thirty-four adult clipped steelhead, twenty-three adult unclipped steelhead, one adult unclipped Chinook, eight juvenile lamprey, five juvenile sculpin, and one adult catfish were returned to the river live and unharmed. In addition, the mortalities included twenty-five juvenile shad, one adult walleye and two adult shad were also placed into the tailrace. In the channel end of the PDS, a stick blockage was discovered once the JCC was dewatered. In the stick blockage is where the mortalities were discovered.

COOLING WATER STRAINERS

Turbine unit cooling water strainers were examined for biologic content once per month from January until June and then again in December. The vast majority of other species found within the strainers were American shad. The number of each group and percent of the individual groups consisted of 1,348 American Shad (53.7%), 1,092 juvenile lamprey (43.6%), 65 salmonid species (<1%), 1 steelhead (<1%), and 1 Siberian prawn (<1%).

Probability of any individual being alive at the time of strainer cleaning was likely more related to time of entry rather than which unit's strainer it was found in. Units 5 and 6 were unable to be inspected because of mechanical issues with the units on December 8. Repairs to unit 5 and 6 will be completed during the unit outages in January and February 2023. Table 19 below reflects the results of this year's main unit cooling water strainer examinations.

Table 16. Cooling Water Strainer Results at Lower Monumental Dam, 2023.

Month	Lamprey Mortality	Live Lamprey	Smolt Mortality	Live Smolts
Jan	1	0	0	0
Feb	11	1	0	0
Mar	48	1	0	0
Apr	100	0	2	0
May	538	14	47	1
Jun	373	1	14	0
Jul	4	0	2	0
Dec	0	0	0	0

INVASIVE SPECIES

During winter maintenance, other dewatering activities and monthly mussel station examinations, no issues were found.

A total of 2,867 Siberian prawns were removed from the sample and disposed of in landfills in accordance with the Washington Department of Fish and Wildlife (WDFW) permit requirements this season. Siberian prawns appeared in many of the samples during the later spring and summer, with the last ones found in the samples during the last few weeks of September.

AVIAN PREDATION

Avian Predation-General

Areas of avian predation monitoring included forebay, turbine discharge, spillway discharge and JFF bypass outfall. Deterrent measures included bird wires across the tailrace of the powerhouse, water cannon sprinklers at the exit of the bypass outfall pipe, bird deterrent spikes at common perching areas, and hazing (April 1 through June 2) under the animal control contract with United States Department of Agriculture-Wildlife Services (USDA-WS). Two shift hazing coverage (daylight to dusk) occurred from May 3 to June 2. After the USDA-WS staff was no longer hazing due to the end of the contract season, JFF staff hazed birds on several days when the trigger point was exceeded. Between July 12 to August 4, JFF staff hazed birds due to the number of piscivorous birds feeding in the tailrace area.

Avian predators tend to rest in the forebay and chase juvenile fish as they jump. They also spend time perched on the lock wall facing the tailrace. At the downstream navigation lock guidewall, bird wires were added along the top rail of the handrail during winter 2008-2009, which effectively reduced the perching previously seen there, however, to a great extent the perching only relocated to the deck in front of the handrails.

The bird observations take place during fish ladder inspections with supplemental counts by WS or JFF personnel on days with no ladder inspection. Figure 3 shows the daily count of gulls, cormorants, grebes and pelicans in the tailrace from April 1 to September 30.

The outfall bird cannon was placed into service on March 17 at 1430 hours. Due to a power outage which occurred on August 18 at 1615 hours, the outfall cannon had to be restarted. Another power outage at the JFF on August 22 at 0655 hours and then again at 1315 hours caused the outfall cannon to lose function. The outfall cannon had to be restarted both times. The outfall bird cannon was winterized on October 13.

A broken anti-bird wire was noticed broken/missing on the September 23 inspection. USDA-WS was contacted, and the repair was made on October 13 by JFF and USDA-WS staff.




Figure 3. Daily count of gulls, cormorant, grebes and pelicans in the tailrace and forebay, 2023.

Adult Fishway Inspection Bird Counts/Wildlife Services Bird Monitoring

These inspections were conducted at random times and contain counts during active bird hazing as well as in its absence. On Mondays and Thursdays (April 1 through June 2) USDA-WS contracted employees collected bird information in the same format as the ladder inspection data and this information was added to the spreadsheet for inclusion in this report. During daylight hours, gulls were present if hazing was not occurring. High juvenile fish numbers passing the dam via spill related to higher gull numbers. In the absence of hazing, gulls appeared to be effective at feeding in the tailrace areas. Each ladder inspection included an avian predator count section for five areas that included forebay (FB), spillway (SWT1), under the bird wires of the turbine discharge (PHT1), downstream of the bird wires below the turbine discharge (PHT2), and lastly the juvenile bypass outfall (JFOF). Each area included counts of both foraging and resting birds. The following summarizes the data collected from April 1 through September 30 of the 2023 season. The averages offered in each category include all data through the period; it is an average of all the fish ladder inspections, USDA-WS and JFF supplemented bird monitoring inspections for that condition (feeding/resting) in each zone.

Gulls

Gull numbers were highest from April 11 to May 31 with later waves from June 4 to 10 after the end of hazing efforts and July 8 until September 30 after the end of barge transport. In all areas, gull numbers dropped after May 31 as juvenile salmonid numbers became increasingly sparse and then numbers increased again as the young of the year American shad became abundant in the area.

The maximum number of gulls observed in all areas was 189 (May 12), with an overall daily average of 34.5 gulls. The number of gulls feeding in the FB ranged from 0 to 25 (May 19) and averaged 1.1 while the number of gulls resting in the FB ranged from 0 to 120 (May 8) and averaged 12.7 gulls. Gulls in the FB

are typically seen resting on the navigation lock guide wall. The number of gulls feeding in SWT1 ranged from 0 to 95 (May 19) and averaged 7.7 while the number of resting gulls in SWT1 ranged from 0 to 75 (September 18) and averaged 7.5. Gulls in SWT1 are typically seen avoiding the pyrotechnics of the hazers firing over the spillway discharge from the navigation lock deck (elevation 536). The number of gulls feeding in PHT1 ranged from 0 to 5 (September 20) and averaged <1 gull. The number of gulls resting in PHT1 ranged from 0 to 6 (September 20) and averaged <1 gull. PHT1 gulls are only typically seen when the hazer is not present. The number of gulls feeding in PHT2 ranged from 0 to 80 (May 20) and averaged 3.3 gulls while the number of resting gulls in PHT2 ranged from 0 to 37 (July 18) and averaged <1 gull. Gulls in PHT2 are only typically seen when the hazer is not present. The number of feeding gulls at the JFOF ranged from 0 to 25 (May 19) and averaged 1.5 gulls. Gulls were not observed resting at the JFOF. JFOF gulls are typically seen when large numbers of juvenile salmonids are bypassed.

Hazing was effective at moving gulls out of the area. Two shifts were used to provide daylight to dusk coverage throughout the historic peak of salmonid outmigration. (Table 20) The second shift of hazing was equally as effective as the morning shift. On days when hazing did not occur, but fish passage numbers were high, the birds returned and resumed normal feeding behaviors. Gull numbers correlated well with the peak of the juvenile fish outmigration this season, as has been the rule in the past. Observers noticed that during higher spring spill volume, large numbers of feeding gulls were seen just below SWT1, so were not included in the tailrace observation numbers. This newly observed feeding location also decreased the effectiveness of hazing efforts.

Cormorants

Overall cormorant abundance was fairly consistent throughout the season. Cormorants tend to be more prevalent in the fall and winter than during juvenile salmonid outmigration. The maximum number of cormorants observed in all areas was 84 (September 29), with an overall daily average of 10.9 cormorants. The number of cormorants feeding in the FB ranged from 0 to 8 (April 21) and averaged <1 cormorants while the number of cormorants resting in the FB ranged from 0 to 17 (August 27) and averaged 2 cormorants. Cormorants in the FB are commonly seen foraging and are impervious to hazing. The number of cormorants feeding in SWT1 ranged from 0 to 49 (September 20) and averaged 1.1 cormorants while the number of resting cormorants in SWT1 ranged from 0 to 76 (September 29) and averaged 4.6 cormorants. Cormorants in SWT1 are not effectively prevented from foraging by the pyrotechnics of the hazers. The number of cormorants feeding in PHT1 ranged from 0 to 15 (September 4) and averaged <1 cormorant while the number of cormorants resting in PHT1 ranged from 0 to 4 (September 17) and averaged <1 cormorants. Cormorant abundance in PHT1 is erratic and individuals are impervious to hazing. The number of cormorants feeding in PHT2 ranged from 0 to 32 (September 5) and averaged 1 cormorant while the number of resting in PHT2 ranged from 0 to 2 (May 4) and averaged 1.7 cormorants. Cormorant observations in PHT2 are similar to those in PHT1. The number of cormorants feeding at the JFOF ranged from 0 to 4 (August 30) and averaged <1 cormorant while the number of cormorants resting ranged from 0 to 1 (April 11) and averaged <1 cormorant. Feeding cormorants are frequently underwater and hard to accurately count

Terns

Tern numbers observed were extremely high when compared to past seasons. The first one was observed on June 30. The maximum number of terns observed in all areas was 27 (July 18), with an overall daily average of 2.3 terns. The number of terns feeding in the FB ranged from 0 to 2 (July 23) and averaged <1 tern while the number of cormorants resting in the FB ranged from 0 to 17 (August 21) and averaged 1 tern. The number of terns feeding in SWT1 ranged from 0 to 11 (July 22) and averaged <1 tern while the number of resting terns in SWT1 ranged from 0 to 12 (July 18) and averaged <1 tern. There were no terns observed feeding or resting in PHT1. The number of terns feeding in PHT2 ranged from 0 to 7 (July 24) and averaged <1 tern while the number of resting in PHT2 ranged from 0 to 4 (July 11) and averaged <1 tern. Terns were not observed feeding or resting at the JFOF

Grebes

Grebes were only seen in the forebay (FB) during the season. The number of grebes feeding in FB ranged from 0 to 8 (June 14) and averaged <1 grebe while no grebes were observed resting in the FB. Grebes are frequently underwater and hard to accurately count.

Pelicans

The first pelican was observed on April 13 with the last observation on September 18. The maximum number of pelicans observed in all areas was 49 (May 2), with an overall daily average of 5 pelicans. The number of pelicans feeding in the FB ranged from 0 to 9 (April 28) and averaged <1 pelican with the number of pelicans resting in the FB ranging from 0 to 7 (April 26) and averaging <1 pelican. Pelicans in the FB are typically seen cruising as a group, generally along the north shoreline. The number of pelicans feeding in SWT1 ranged from 0 to 37 (May 18) and averaged 1.6 pelicans while the number of pelicans resting in SWT1 ranged from 0 to 3 (April 20) and averaged <1 pelican. Pelicans in SWT1 are typically not impacted by WS activities. The number of pelicans feeding in PHT1 ranged from 0 to 11 (July 21) and averaged <1 while the number of pelicans resting in PHT1 ranged from 0 to 4 (July 16) and averaged <1 pelican. The number of pelicans feeding in PHT2 ranged from 0 to 34 (May 26) and averaged 1.8 pelicans while the number of pelicans resting in PHT2 ranged from 0 to 7 (June 4) and averaged <1 pelican. The number of pelicans feeding at the JFOF ranged from 0 to 6 (May 19) and averaged <1 pelican while the number of pelicans resting at JFOF ranged from 0 to 2 (July 20) and averaged <1 pelican. Pelicans near the JFOF are typically seen when large numbers of juvenile salmonids are bypassed.

Avian Hazing-United States Department of Agriculture-Wildlife Services

Deck hazers worked eight-hour shifts for seven days a week. Start and end times varied to reduce habituation of birds. Bird hazing efforts by USDA-WS personnel began on April 1. The USDA-WS hazing program is outlined in Table 20.

Table 17. USDA-WS hazing program schedule for Lower Monumental Dam, 2023

Personnel	Days	Dates	Shift
WS Hazer #1	Monday–Friday	4/1/2023–6/2/2023	Regular Coverage
WS Hazer #2	Monday–Friday	5/3/2023–6/2/2023	Peak Season
WS Hazer #3	Saturday & Sunday	4/1/2023–6/2/2023	Regular Coverage
WS Hazer #4	Saturday & Sunday	5/3/2023–6/2/2023	Peak Coverage

RECOMMENDATIONS

Research converting the pipe system between the PIT facility counter tanks and the PIT facility holding tank exits with an open system that eliminates the need to hold fish in the PIT system holding tanks.

Research a new juvenile count system to replace the Smith-Root count system.

Remove sand and debris from the supply conduits and replace all original ladder diffuser grates, support structures, and mud valves.

Replace staff gauges with fiberglass reinforced plastic staff gauges.